

# Local Therapies for Localised Neuropathic Pain

Arun Bhaskar\*, Consultant in Pain Medicine & Anaesthesia

Rahul Mittal\*, Specialist Trainee in Clinical Oncology

*\*The Christie NHS Foundation Trust, Withington, Manchester*

## Introduction

Localised neuropathic pain commonly arises due to damage to a peripheral nerve; but can also arise from damage to or hyper-reactivity of nerve plexus, nerve root, or occasionally from central processes<sup>1</sup>. This type of pain can cause significant negative impact on quality of life. Systemic drugs used to treat Neuropathic pain, such as antidepressants, anticonvulsants and opioid analgesics can also cause significant side effects such as sedation, drowsiness and cognitive dysfunction that increase difficulties for the patient<sup>2</sup>.

This paper will address local treatment options that are effective for localised neuropathic pain whilst avoiding the long-term systemic side effects of medications; it should be mentioned that there are well known complications that could arise after invasive therapies, but these are minimised in the hands of experienced clinicians and by using advanced imaging techniques and other precautionary measures.

## Topical Pharmacological Agents

Amitriptyline and Gabapentin are well known systemic drugs used in the management of neuropathic pain. However, they are also used as topical agents, apparently with some success, particularly the use of gabapentin in vulvodynia. The mechanisms for topical relief of pain seem to differ from their central effects, and the mechanism of action is unknown. The uses of many agents are limited to sympathetically mediated pain<sup>3</sup> and treatment-resistant pain, as they are not freely available, although there are ongoing studies looking into the use of various combinations of different agents topically.

Amitriptyline cream has been used in the management of post-traumatic neuropathic pain<sup>4</sup>, but in studies when compared with 5% lidocaine cream<sup>5</sup> and 3.3% doxepin cream<sup>6</sup>, amitriptyline cream was found to be less efficacious. However, when combined with ketamine cream, topical amitriptyline showed promising results in neuropathic pain states<sup>7</sup> and also in the management of refractory proctodynia<sup>8</sup>. The use of the above combined with Baclofen in a cream form has been shown to be effective in chemotherapy-induced peripheral neuropathy<sup>9</sup>. Topical Gabapentin has been used in the management of localised and generalised vulvodynia<sup>10</sup> and anecdotally has been tried in other neuropathic pain states. Lidocaine creams and EMLA are also used, especially for mucous membranes, but has largely been replaced by 5% Lidocaine plasters on skin, which is discussed below.

Ketamine is used as a systemic agent in the management of neuropathic pain, but its use as a topical gel has also been explored<sup>11</sup>. It has been used in the management of oral mucositis<sup>12</sup> and its use in combination with amitriptyline cream has been discussed before<sup>7,8,9</sup>. Clonidine is another agent that has been tried topically<sup>13</sup> and it works by blocking the emerging pain signals at peripheral terminals via alpha-2 adrenoceptors without producing undesirable central side effects<sup>14</sup>. GTN spray was also tried with some success in the management of painful diabetic neuropathy either as a sole agent or in combination with anticonvulsants<sup>15</sup>.

The agent that is commonly used for localised neuropathic pains including painful diabetic neuropathy and post-herpetic neuralgia is capsaicin<sup>16,17</sup>. It is usually used as a 0.025% or 0.075% cream and is licensed for the management of painful diabetic neuropathy<sup>18,19</sup>. Combination of doxepin and capsaicin cream was found to produce faster results in another study<sup>20</sup>. Some patients find it difficult to be compliant with the treatment due to the burning nature of the cream and it needs to be applied 3–4 times a day for several weeks before producing pain relief. The 8% capsaicin patch, which is a single application could improve on this and is discussed in detail below.

## 5% Lidocaine Plaster

Topical 5% lidocaine plaster has been around since late 1999 and is a topical analgesic, which is recommended by some algorithms as first line therapy for the treatment of localised, peripheral neuropathic pain<sup>21,22</sup>. Each 10 cm x 14 cm plaster contains 700 mg (5%w/w) lidocaine in a white hydrogel plaster containing adhesive material and is applied for a period no longer than 12 hours with a subsequent plaster-free interval for another 12 hours<sup>23</sup>; a maximum of three plasters are used at a time<sup>23</sup>. The application of the plaster over the intact skin of the affected area is usually well-tolerated and application site side effects like erythema, rash and pruritus are mostly self-limiting<sup>24</sup>. Patients need to be followed up after 2–4 weeks and if there is no analgesic benefit in that period, it is unlikely to get more benefit than the barrier effect.

The mechanism of action is still unclear, but the local analgesic effect is by stabilising neuronal membranes, and from animal studies it is thought to down-regulate the sodium channels present in the peripheral nerve endings, reducing ectopic nociceptive pain signal transmission<sup>25</sup>. As a long-term consequence, reduction of peripheral nerve input may counteract central sensitisation. Lidocaine induces a TRPV1 dependent release of CGRP, which is a key component of neurogenic inflammation and long-term topical application



of lidocaine reduces nerve fibre density in the epidermis<sup>26,27</sup>; this degeneration may explain the observed loss of pain<sup>28</sup>. Additionally, the plaster acts as a barrier protecting the skin from mechanical stimuli, which can trigger pain sensations due to allodynia and hyperpathia.

The current licensed indication for 5% lidocaine plaster is for post-herpetic neuralgia<sup>23</sup>. Various studies evaluating the long-term efficacy, safety and tolerability of the lidocaine plaster show consistent and sustained long-term pain relief for up to 24 months and longer<sup>29</sup>. The data further confirms that in newly diagnosed patients, pain relief was achieved at 1 week of treatment and subsequent reductions occurred in the weeks thereafter. 5% lidocaine plasters have also been successfully used in patients with other types of neuropathic pain states, either as a single agent<sup>30</sup> or in combination<sup>31</sup>. Promising results have also been obtained with the use of 5% lidocaine patch for the treatment of mild-to-moderate carpal tunnel syndrome, the efficacy being comparable to that of the more invasive anaesthetic/corticosteroid injection otherwise used<sup>32</sup>. It was found that the plaster was well tolerated by the patients who also reported a low side effect profile.

5% lidocaine plasters are also used in treating other pain conditions including myofascial pain syndrome (MPS), caused by trigger points in muscle or muscle fascia. The existing standard treatment for this is infiltration of the trigger points with local anaesthetic +/- steroids; however this intervention normally can produce some discomfort in the patients. The application of topical 5% lidocaine plasters for the symptoms of MPS was found to be effective and highly acceptable to patients with MPS<sup>33</sup>. The use of 5% lidocaine plasters in three different chronic pain conditions for two weeks also had a significant reduction and impact on all pain measures assessed by neuropathic pain scoring<sup>34</sup>. It has also been shown that 5% lidocaine plaster can be used as an additional therapeutic option in chronic back pain<sup>35</sup> and also showed no statistically significant difference between using the 5% lidocaine plaster versus COX-2 inhibitors in treating osteoarthritis pain<sup>36</sup>.

In summary, 5% lidocaine plaster is an effective means of treating ongoing pain and allodynia due to localised neuropathic pain and diverse peripheral neuropathies. The sustained efficacy, combined with its safety and tolerability profile results in a favourable benefit-risk ratio, suggesting its safety for long-term use. However, it is to be noted that, being licensed for post-herpetic neuralgia, the predominant use is still in neuropathic pain and not for musculoskeletal pain.

### 8% Capsaicin Patch

Capsaicin (8-methy-N-vanillyl-6-nonenamide) is the active component of the chilli pepper and is an agonist of the transient receptor potential vanilloid-1 receptor (TRPV1)<sup>37</sup>. Topical capsaicin creams, 0.025% and 0.075%, have been used over twenty years in the management of peripheral neuropathic pains<sup>38</sup>, but tend to have limited analgesic effect, as it needs to be applied up to four times

daily for several weeks, which could lead to reduced compliance rates. Some patients also complain of experiencing a burning sensation when using the cream<sup>39</sup>. The low dose topical capsaicin cream may reduce neuropathic pain in PHN but in one trial it failed show efficacy in HIV-associated distal sensory polyneuropathy<sup>40</sup>.

A high concentration capsaicin (8%w/w) patch was approved in the UK in 2010 for the treatment of PHN and other peripheral neuropathic states other than diabetic neuropathy. Each 14 cm x 20 cm patch, with a total of 179 mg of capsaicin (640 mcg/cm<sup>2</sup>) in the silicone-adhesive mixture, is designed to deliver a therapeutic dose of capsaicin during a single application of 30-60 minutes. Capsaicin causes an initial enhanced stimulation of TRPV1-expressing cutaneous nociceptors that may be associated with painful sensations, followed by a reduction in the TRPV1-expressing nociceptors, leading to prolonged analgesia<sup>41</sup>. Over a period of months, there may be a gradual re-emergence of the neuropathy due to nerve fibre re-innervation of the treated area. The mechanism of action is thought to be selective and reversible defunctioning of cutaneous sensory nerve endings expressing TRPV1. However, large myelinated nerve fibres are unaffected, thereby preserving the integrity of protective touch, vibratory and thermal connections<sup>41</sup>. This is important as it means capsaicin treatment can lead to reduction in pain without clinically relevant changes in protective sensation.

Evidence from several controlled clinical trials have demonstrated that applying a single 60 minute high concentration capsaicin patch can significantly reduce neuropathic pain in patients with PHN<sup>42-45</sup>. The studies also confirmed that it could be effective as a monotherapy or in conjunction with other pain analgesia<sup>40,42,43</sup> and more importantly, the pain relief was maintained for up to at least 12 weeks<sup>43-46</sup>. The patch showed greater improvement in mean pain scores at all assessment times<sup>42-46</sup>. A randomised double blind controlled study in patients with PHN for at least three months showed that following treatment with the 8% capsaicin patch, pain scores began declining as early as the end of day 1 and were on average reduced by 36% during the first week<sup>46-47</sup>.

Studies have also shown that the 8% capsaicin patch is effective in HIV-associated distal sensory polyneuropathy, a painful condition with limited effective treatments<sup>40,48,49</sup>. A multicentric double blind controlled dose-finding study, which evaluated the safety and efficacy of the 8% patch but studied with different durations of application - 30, 60 or 90 minutes in treating PHN<sup>46</sup>. In summary the authors found that a single topical application of the patch provided pain reduction for up to 3 months<sup>46</sup>. Treatment duration of 60 minutes was found to be the lowest effective dose time and generally well tolerated, but for the treatment of painful HIV neuropathy of the feet, it was found that a 30 minutes application gave similar analgesic effect for a similar duration<sup>48-49</sup>.

Side effects of 8% capsaicin are predominantly localised, such as application site pain, pruritus and papules; these responded to localised cooling and analgesics, and lasted only for a few days<sup>45,50</sup>. A transient, but clinically insignificant rise in blood pressure has also been noted during the duration of the patch application and was



self-limiting<sup>45</sup>. Exposure to capsaicin may result in nasopharyngitis and this can be avoided by careful removal, rolling the patch inwards to avoid contact, for safe disposal.

In summary, high dose capsaicin patch is generally well tolerated by patients and appears to have some potential advantages over some of the current first line agents in term of side effect profile. Poor compliance from patients for capsaicin cream is avoided, as the whole therapeutic dose is delivered during a single supervised application.

### **Percutaneous Electrical Nerve Stimulation Therapy (PENS)**

PENS therapy is a type of minimally invasive peripheral nerve stimulation used in the treatment of chronic peripheral neuropathic pain. Since the advent of the “Gate Control Theory” of pain<sup>51</sup>, electrical stimulation has been used to provide analgesia. Pain due to nociceptive stimuli is carried via C-fibres to the dorsal horn at laminae 1 & 2 and then onwards via the spinothalamic tract to terminate in the sensory cortex. Transcutaneous Electrical Nerve Stimulation (TENS) causes a reduction of pain by stimulating peripheral inhibitory fibres. This may not always be effective in some neuropathic pain states, as C-fibre termination at laminae 1 & 2 ceases as a result of peripheral nerve damage and is replaced by A $\beta$ -fibres at the lamina<sup>52</sup>. The A $\beta$ -fibres projecting into the lamina can cause an exaggerated nociceptive response to what are normally innocuous stimuli<sup>53</sup> and hence these patients may not tolerate TENS, but may be more likely to benefit from PENS therapy. PENS uses a frequency-dependent electrical current (usually 2 Hz and 100 Hz alternating every 3 seconds), to modulate the activity of peripheral nerves and can be used to stimulate a single peripheral nerve<sup>54</sup> or the most peripheral branches of a peripheral nerve as field stimulation<sup>55</sup>. The electrical current is delivered through a 21 gauge disposable probe of varying lengths of 20 mm to 200 mm inserted percutaneously to target the affected area for a period of 25 minutes.

PENS therapy offers patients the potential for intermediate relief and ongoing management of chronic peripheral neuropathic pain conditions like scar pain including post-thoracotomy, post-mastectomy and post-hernia repair pains<sup>56</sup>. It has also been used with success in intractable cluster headaches<sup>57</sup> and occipital neuralgias<sup>58</sup>. PENS gives good analgesic results in patients with post-irradiation allodynia and following trigeminal field neuropathy following head and neck cancer surgery and radiotherapy<sup>59</sup>. These patients reported improved quality of life and could make a significant reduction in opioid analgesia and also systemic neuropathic pain agents<sup>60</sup>.

PENS therapy is contraindicated in the presence of demand type cardiac pacemakers and patient refusal. Caution is to be exercised in patients known to have heart disease, epilepsy, on anticoagulants and in pregnancy. PENS therapy is best deferred when there is local or systemic infection or coagulopathy. Possible adverse events may include temporary flare of pain, bleeding or haematoma at probe insertion site, and the possibility of infection at the probe entry point; misplacement of the probe may cause temporary neuropraxia, though this is very rare.

PENS can be used as a diagnostic therapy to identify patients who are likely to benefit from a permanently implanted neurostimulator and could reduce the incidence of patients who fail to benefit from a lengthy, potentially traumatic and costly procedure<sup>61</sup>. Distinguishing patients who may stand to benefit from permanently implantable neurostimulation devices, from those who may not, and the ability to recognise those who will gain an enhanced quality of life through ongoing PENS therapy, offers clear clinical and economic advantages when considered in conjunction with other neurostimulation therapies. It has also been used with success in patients with focal neuropathic pain in patients with cancer who require frequent MRI scans which make an implantable neurostimulator contraindicated<sup>60</sup>.

PENS offers the option of a minimally invasive neuromodulation therapy that can be carried out in a procedure room, either as a diagnostic tool prior to implantable stimulators or a continuing treatment option for localised neuropathic pain states. It is associated with a low risk profile and is a cost-effective option when pharmacological options are either ineffective or when the effective dose causes unacceptable side effects to the patient that could significantly impact on their quality of life. The evidence base for PENS treatment is growing, but are currently limited to case series and case reports; there needs to be carefully conducted randomised controlled clinical trials (RCTs) to further validate and provide hard evidence of therapeutic efficacy.

### **Radiofrequency Ablation**

Radiofrequency techniques have been in use for more than 30 years in the management of chronic pain conditions<sup>62</sup>, particularly back<sup>63</sup> and neck pain and also neuropathic conditions like trigeminal neuralgia and other refractory neuropathic pain states<sup>64</sup>. There is a wealth of clinical experience and success in the use of both conventional (thermal) and pulsed radiofrequency techniques, including several large case series, but there are only a few high quality randomised controlled trials to validate these results<sup>65</sup>. However, it is widely used in clinical practice in the management of focal neuropathic pain<sup>66</sup>.

The principles and procedure of radiofrequency have been well established and both conventional and pulsed lesioning are used in common practice<sup>67,68</sup>. Conventional radiofrequency using a thermal lesion at temperatures higher than 65 degrees Celsius can produce significant damage to neural tissue; pulsed lesioning is done at lower temperatures around 42 degrees Celsius and thus gives rise to minimal damage and hence a lower incidence of complications<sup>69</sup>. Experimental studies have shown that though no gross change is seen on both treatments, electron microscopic examination has revealed minimal disruption of the myelin envelope of the axon<sup>70</sup>, indicating that the analgesic effect of radiofrequency technique is not just by causing a thermal lesion. Pulsed radiofrequency defines an electromagnetic field in the vicinity of a lesion, modifying neuro-cellular function with minimal cellular destruction<sup>71</sup>. The mechanisms of the analgesic action by pulsed radiofrequency by the enhancement of descending nor-adrenergic and serotonergic



systems<sup>72</sup> have also been described in animal studies. There may also be an anti-nociceptive effect produced simply due to radiofrequency application<sup>73</sup>.

The use of radiofrequency lesioning for trigeminal neuralgia has been in practice for nearly forty years<sup>74</sup>; the technique is well described<sup>75</sup> and studies have shown that conventional radiofrequency is far more effective in sustaining analgesia than pulsed lesioning<sup>76</sup>. Percutaneous radiofrequency rhizotomy has been effective in idiopathic trigeminal neuralgia, symptomatic facial pain and herpes zoster of the fifth cranial nerve, but not effective in atypical facial pain<sup>77</sup>. It has been the recommended option for treating trigeminal pain in the elderly, but microvascular decompression remains the first line choice for younger patients<sup>78</sup>. The procedure is not without complications and apart from pain and bruising, major complications such as other cranial nerve palsies<sup>79</sup>, subarachnoid haemorrhage<sup>80</sup> and rhinorrhea<sup>81</sup> have also been reported. Targeting the sphenopalatine ganglion<sup>82</sup>, particularly using advanced imaging including CT-guidance<sup>83</sup> and virtual reality techniques<sup>84</sup> has higher efficacy and reduced risk of complications. Radiofrequency neurolysis of the infra-orbital nerve has been done when trigeminal ganglion or posterior root ablation is contraindicated<sup>85</sup>. Pulsed radiofrequency has been used in the treatment of occipital neuralgia with the pain improving substantially in more than half the number of patients<sup>86,87</sup>.

Several other neuropathic pain conditions have been treated with radiofrequency with good to exceptional results as reported in case series. Pulsed radiofrequency of the lateral cutaneous nerve of the thigh has been used in successfully treating meralgia paraesthetica<sup>88</sup>. It has also been used with success in managing groin pain and orchialgia<sup>89</sup> and also refractory pudendal neuralgia<sup>90</sup>. There are case reports of effective radiofrequency lesioning for obturator nerve neuropathy<sup>91</sup> for conditions such as hip arthritis. The effects of producing a radiofrequency lesion adjacent to the dorsal root ganglion in patients with thoracic segmental pain has also been reported, with 67% of patients reporting pain relief at 8 weeks and half the patients having relief at 36 weeks<sup>92</sup>. More recently it has been shown that pulsed radiofrequency of the dorsal root ganglion is superior to pharmacotherapy or pulsed radiofrequency of the intercostal nerves in the treatment of chronic postsurgical thoracic pain<sup>93</sup>. There are also case reports of CT-guided dorsal root ganglion thermocoagulation for treating post herpetic neuralgia, where the majority of patients had excellent pain relief and others had a decrease in symptoms<sup>94</sup>. Radiofrequency lesioning of the thoracic paravertebral nerve for refractory neuropathic pain following breast cancer surgery has also been reported<sup>95</sup>.

There are several case reports and case series regarding the use of radiofrequency ablation in the management of lower limb phantom pain, including pulsed radiofrequency of the sciatic nerve<sup>96</sup>. The patients reported improved prosthetic tolerance and reduction of oral analgesic medications. The analgesic effect is thought to be due to inhibition of evoked synaptic activity<sup>97</sup>. It has also been reported that there could be increased pain, new pain syndromes and also abnormal sweating in the limb stump<sup>98</sup>. Pulsed radiofrequency has also been used in treating myofascial trigger points and scar

neuromas with 67% of the patients reporting pain relief lasting between 6 to 12 weeks<sup>99</sup>. It has also been used in treating patients with chronic neuropathic spinal pain after failed conservative management<sup>100</sup>. A 12-year long-term retrospective analysis of using radiofrequency application to areas of pain in the treatment of neurogenic heel pain has shown a high success rate with few associated risks and less post-operative morbidity<sup>101</sup>. Radiofrequency ablation has been used with success for the management of prolonged moderate to severe heel pain associated with plantar fasciitis<sup>102</sup>.

Cervico-thoracic or lumbar sympathectomy for neuropathic pain and complex regional pain syndrome (CRPS) has poor evidence in literature due to lack of large well-designed studies<sup>103</sup>. However, radiofrequency lesions of the stellate ganglion have been used with success in CRPS Type 2, ischaemic pain, cervicobrachialgia and post-thoracotomy pain<sup>104</sup>. Pulsed radiofrequency lumbar sympathectomy has also been used in the management of CRPS in a patient with spinal injury<sup>105</sup>. However, the evidence base for the use of radiofrequency ablation for these conditions has not yet been validated by randomised controlled trials<sup>106</sup>.

## Summary

Topical agents such as 5% Lidocaine plasters, 8% Capsaicin patch and procedures such as PENS Therapy give the added option of delivering pain relief locally without the side-effects of systemic drugs for localised neuropathic pain. Though radiofrequency techniques are widely used and there is moderate evidence for facet joint neurotomy, more well-designed studies need to be conducted to gather more evidence in its use in other neuropathic pain states.

## REFERENCES

1. Woolf C. Pain: moving from symptom control toward mechanism-specific pharmacologic management. *Ann Intern Med.* 2004;140:441-451.
2. Beydoun A, et al. Mechanistic stratification of antineuralgic agents. *J Pain Symptom Manage.* 2003;25(suppl 5):S18-30.
3. Knotkova H, Pappagallo M. Adjuvant analgesics. *Med Clin North Am.* 2007 Jan;91(1):113-24
4. Liebergts R, Kopsky DJ, Keppel Hesselink JM. Topical amitriptyline in post-traumatic neuropathic pain. *J Pain Symptom Manage.* 2011 Apr;41(4):e6-7
5. Ho KY, Huh BK, White WD, Yeh CC, Miller EJ. Topical Amitriptyline versus lidocaine in the treatment of neuropathic pain. *Clin J Pain* 2008 Jan 24(1):51-55



6. Besson M, Desmeules J, Piguet V. What is the place of topical analgesia in neuropathic pain? *Rev Med Suisse* 208 Jun 18;4(162): 1498, 1500, 1502-4
7. Lynch ME, Clark AJ, Sawynok J, Sullivan MJ. Topical 2% amitriptyline and 1% ketamine in neuropathic pain syndromes; a randomized, double-blind, placebo-controlled trial. *Anesthesiology*. 2005 Jul; 103(1):140-6
8. Lehman JS, Sciallis GF. Effective use of topical amitriptyline hydrochloride 2.5% and ketamine hydrochloride 0.5% for analgesia in refractory proctodynia. *J Drugs Dermatol*. 2008 Sep;7(9)0: 887-9.
9. Barton DL, Wos EJ, Qin R, Mattar BI, Green NB, Lanier KS, Bearden JD 3rd, Kugler JW, Hoff KL, Reddy PS, Rowland KM Jr, Riepl M, Christensen B, Lopirizzi CL. A double-blind, placebo-controlled trial of a topical treatment for chemotherapy-induced peripheral neuropathy: NCCTG trial N06CA. *Support Care Cancer* 2011 Jun;19(6):833-41. *Epub* 2010 May 25.
10. Boardman LA, Cooper AS, Blais LR, Raker CA. Topical gabapentin in the treatment of localized and generalized vulvodynia *Obstet Gynecol*. 2008 Sep; 112(3):579-85
11. Gammaitoni A, Gallagher RM, Welz-Bosna M. Topical ketamine gel: possible role in treating neuropathic pain. *Pain Med*. 2000 Mar;1(1):97-100.
12. Slatkin NE, Rhiner M. Topical ketamine in the treatment of mucositis pain. *Pain Med*. 2003 Sep;4(3):298-303
13. Motsch J, Kamler M. Alpha 2-adrenergic agonists. Use in chronic pain – a meta-analysis *Schmerz*. 1997 Oct 24;11(%):339-344.
14. Dogrul A, Uzbay IT. Topical Clonidine antinociception. *Pain* 2004 Oct;111(3):385-91
15. Agrawal RP, Goswami J, Jain S, Kochar DK. Management of diabetic neuropathy by sodium valproate and glyceryl trinitrate spray: a prospective double-blind randomized placebo-controlled study. *Diabetes Res Clin Pract*. 2009 Mar;83(3):371-8.
16. Derry S, Lloyd R, Moore RA, McQuay HJ. Topical capsaicin for chronic neuropathic pain in adults. *Cochrane Database Syst Rev*. 2009 Oct 7;(4):CD007393. *Review*.
17. Mason L, Moore RA, Derry S, Edward JE, McQuay HJ. Systematic review of topical capsaicin for the treatment of chronic pain. *BMJ*. 2004 Apr 24;328(7446):991. *Epub* 2004 Mar 19. *Review*.
18. Murad MH, Smith SA. Review: TCAs, anticonvulsants, opioids and capsaicin cream are effective for diabetic neuropathy. *Evid Based Med*. 2008 Feb;13(1):21.
19. Jensen PG, Larson JR. Management of painful diabetic neuropathy. *Drugs Aging*. 2001; 18(10):737-49. *Review*
20. McClean G. Topical application of doxepin hydrochloride, capsaicin and a combination of both produces analgesia in chronic human neuropathic pain: a randomized, double-blind, placebo-controlled study. *Br J Clin Pharmacol*. 2000 Jun;49(6):574-9
21. Finnerup NB, Otto M, McQuay HJ, Jensen TS, Sindrup SH. Algorithm for neuropathic pain treatment: an evidence based proposal. *Pain*. 2005 Dec 5;118(3):289-305. *Epub* 2005 Oct 6.
22. Dworkin RH, O'Connor AB, Backonja MM et al. Treatment of neuropathic pain: an overview of recent guidelines. *Am J Med*. 2009 Oct;122(10 Suppl):S22-32. *Review*.
23. Grunenthal Ltd. Versatis 5% Lidocaine plaster: summary of product characteristics (SPC)
24. Meier T, Wasner G, Faust M, Kuntzer T, Ochsner F, Hueppe M, Bogousslavsky J, Baron R. Efficacy of lidocaine patch 5% in the treatment of focal peripheral neuropathic pain syndromes: a randomized, double blind, placebo-controlled study. *Pain*. 2003 Nov;106(1-2):151-8.
25. Nicolaou A, Nicholson B, Hans G, Brasseur L. Outcome predictors for treatment success with 5% lidocaine medicated plaster in low back pain with neuropathic components and neuropathic pain after surgical and nonsurgical trauma. *J Pain Research*. 2011 Jan 11;4:25-38.
26. Wehrfritz A, Leffler A, Namer B, Müller C, Koppert W. Topical lidocaine in a human pain model reduces pain sensation and quantity of epidermal nerve fibres. *Poster 425: EFIC-Congress* 2009
27. Wehrfritz A, Namer B, Ihmsen H, Müller C, Filitz J, Koppert W, Leffler A. Differential effects on sensory functions and measures of epidermal nerve fiber density after application of a lidocaine patch (5%) on healthy human skin
28. Wasner G, Kleinert A, Binder A, Schattschneider, Baron R. Post-herpetic neuralgia: Topical lidocaine is effective in nociceptor-deprived skin. *J Neurol* (2005) 252:677-686 DOI 10.1007/s00415-005-0717-z
29. Baron R, Mayoral V, Leijon G, Binder A, Steigerwald I, Serpell M. 5% lidocaine medicated plaster versus pregabalin in post-herpetic neuralgia and diabetic polyneuropathy: an open-label, non-inferiority two-stage RCT study. *Curr Med Res Opin*. 2009 Jul;25(7):1663-76.

30. Baron R, Mayoral V, Leijon G, Binder A, Steigerwald I, Serpell M. Efficacy and safety of combination therapy with 5% lidocaine medicated plaster and pregabalin in post-herpetic neuralgia and diabetic polyneuropathy. *Curr Med Res Opin.* 2009 Jul;25(7):1677-87.
31. Hans G, Sabatowski R, Binder A, Boesl I, Rogers P, Baron R. Efficacy and tolerability of a 5% lidocaine medicated plaster for the topical treatment of post-herpetic neuralgia: results of a long-term study. *Curr Med Res Opin.* 2009 May;25(5):1295-305.
32. Nalamachu S, Crockett RS, Mathur DJ. Lidocaine patch 5 for carpal tunnel syndrome: how it compares with injections: a pilot study. *J Fam Pract.* 2006 Mar;55(3):209-14.
33. Affaitati G, Fabrizio A, Savini A, Lerza R, Tafuri E, Costantini R, Lapenna D, Giamberardino MA. A randomized, controlled study comparing a lidocaine patch, a placebo patch, and anesthetic injection for treatment of trigger points in patients with myofascial pain syndrome: evaluation of pain and somatic pain thresholds. *Clin Ther.* 2009 Apr;31(4):705-20.
34. Gammaitoni AR, Galer BS, Onawola R, Jensen MP, Argoff CE. Lidocaine patch 5% and its positive impact on pain qualities in osteoarthritis: results of a pilot 2-week, open-label study using the Neuropathic Pain Scale. *Curr Med Res Opin.* 2004;20 Suppl 2:S13-9.
35. Baliki MN, Geha PY, Jabakhanji R, Harden N, Schnitzer TJ, Apkarian AV. A preliminary fMRI study of analgesic treatment in chronic back pain and knee osteoarthritis. *Mol Pain.* 2008 Oct 25;4:47.
36. Kivitz A, Fairfax M, Sheldon EA, Xiang Q, Jones BA, Gammaitoni AR, Gould EM. Comparison of the effectiveness and tolerability of lidocaine patch 5% versus celecoxib for osteoarthritis-related knee pain: post hoc analysis of a 12 week, prospective, randomized, active-controlled, open-label, parallel-group trial in adults. *Clin Ther.* 2008 Dec;30(12):2366-77.
37. Noto C, Pappagallo M, Szallasi A. NGX-4010. A high concentration capsaicin dermal patch for lasting relief of peripheral neuropathic pain. *Curr Opin Investig Drugs.* 2009 Jul;10(7):702-10.
38. Capsaicin Study Group. Treatment of painful diabetic neuropathy with topical capsaicin: a multicenter, double-blind, vehicle-controlled study. *Arch Intern Med* 151:2225-29, 1991.
39. Capsaicin Study Group. Effect of treatment with capsaicin on daily activities of patients with painful diabetic neuropathy. *Diabetes Care* 15:159-65, 1992.
38. Simpson DM, Brown S, Tobias J. NGX-4010 C107 Study Group. Controlled trial of high-concentration capsaicin patch for treatment of painful HIV neuropathy. *Neurology.* 2008 Jun 10;70(24):2305-13.
41. Kennedy WR, Vanhove GF, Lu SP, Tobias J, Bley KR, Walk D, Wendelschafer-Crabb G, Simone DA, Selim MM. A randomized, controlled, open-label study of the long-term effects of NGX-4010, a high-concentration capsaicin patch, on epidermal nerve fiber density and sensory function in healthy volunteers. *J Pain.* 2010 Jun;11(6):579-87. Epub 2010 Apr 18.
42. Backonja M, Wallace MS, Blonsky ER, Cutler BJ, Malan P Jr, Rauck R, Tobias J; NGX-4010 C116 Study Group. A high-concentration capsaicin patch, for the treatment of postherpetic neuralgia: a randomised, double-blind study. *Lancet Neurol.* 2008 Dec;7(12):1106-12. Epub 2008 Oct 30. Erratum in: *Lancet Neurol.* 2009 Jan;8(1):31.
43. Backonja MM, Malan TP, Vanhove GF, Tobias JK; C102/106 Study Group. NGX-4010. A high-concentration capsaicin patch, for the treatment of postherpetic neuralgia: a randomized, double-blind, controlled study with an open-label extension. *Pain Med.* 2010 Apr;11(4):600-8. Epub 2010 Jan 22.
44. Backonja MM et al. High-concentration capsaicin for the treatment of post-herpetic neuralgia and other types of peripheral neuropathic pain *European Journal of Pain Supplements* Volume 4, Issue 2, August 2010, Pages 170-174
45. Irving GA, Backonja MM, Duntzman E, Blonsky ER, Vanhove GF, Lu SP, Tobias J. NGX-4010 C117 Study Group. A multicenter, randomized, double-blind, controlled study of NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia. *Pain Med.* 2011 Jan;12(1):99-109. doi: 10.1111/j.1526-4637.2010.01004.x. Epub 2010 Nov 18.
46. Webster LR, Tark M, Rauck R, Tobias JK, Vanhove GF. Effect of duration of postherpetic neuralgia on efficacy analyses in a multicenter, randomized, controlled study of NGX-4010, an 8% capsaicin patch evaluated for the treatment of post-herpetic neuralgia. *BMC Neurol.* 2010 Oct 11;10:92.
47. Webster LR, Malan TP, Tuchman MM, Mollen MD, Tobias JK, Vanhove GF. A multicenter, randomized, double-blind, controlled dose finding study of NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia. *J Pain.* 2010 Oct;11(10):972-82. Epub 2010 Jul 24.
48. Simpson DM, Brown S, Tobias J; NGX-4010 C107 Study Group. Controlled trial of high-concentration capsaicin patch for treatment of painful HIV neuropathy. *Neurology.* 2008 Jun 10;70(24):2305-13.



49. Simpson DM, Estanislao L, Brown SJ, Sampson J. An open-label pilot study of high-concentration capsaicin patch in painful HIV neuropathy. *J Pain Symptom Manage.* 2008 Mar; 35(3): 299-306. Epub 2007 Oct 23.
50. Simpson DM, Gazda S, Brown S, Webster LR, Lu SP, Tobias JK, Vanhove GF; NGX-4010 C118 Study Group. Long-term safety of NGX-4010: A high concentration capsaicin patch in patients with peripheral neuropathic pain. *J Pain Symptom Manage.* 2010 Jun;39(6):1053-64.
51. Melzack R, Wall PD. Pain mechanisms: a new theory. *Science.* 1965;150(699):971-9. PMID 5320816.
52. Gracely, R.H., Lynch, S.A., and Bennett, G.J. (1992). Painful neuropathy: altered central processing maintained dynamically by peripheral input. *Pain*, 175-194.
53. Han, J.S. and Wang, Q. (1992). Mobilization of specific neuropeptides by peripheral stimulation of identified frequencies. *Neuro Physiological Science* 7, 176-180.
54. Bhaskar A, Chatha H. Proximal application of Percutaneous Electrical Nerve Stimulation (PENS) for treating neuropathic thigh and knee pain following radical trachelectomy and laparoscopic pelvic lymph node dissection surgery. *Proceeding of 3rd Neuropathic Pain Congress (NeuPSIG)*, Athens 2010.
55. Hu P, Lee M, O'Keeffe D, Tracey I: Peripheral electrical stimulation in Capsaicin induced secondary hyperalgesia: A psychophysical evaluation. *Proceeding of IASP Glasgow*, 2008
56. Raheem T, Raphael J, Duarte R, Akinwunmi J, Cooper R, Bennett A: A multi-centre open prospective study of percutaneous electrical neuro- stimulation in post-surgical scar pain. *Proceedings of ESRA annual congress*, Genoa, Italy, September 2008.
57. Burns B, Watkins L, Goadsby P: Treatment of medically intractable cluster headache by occipital nerve stimulation: long-term follow-up of eight patients. *The Lancet*, March 8 2007, DOI:10.1016/S0140- 6736(07)60328-6.
58. Magis D, Allena M, Bolla M, De Pasqua V, Remacle J, Schoenen J: Occipital nerve stimulation for drug- resistant chronic cluster headache: a prospective pilot study. *The Lancet*, March 8 2007, DOU:10.1016.S1474- 4422(07)70058-3.
59. Bhaskar A, White A, Patel K. Percutaneous Electrical Nerve Stimulation (PENS) Therapy for the management of focal neuropathic pain – a case series of 42 cancer patients. *Proceeding of 10th INS World Congress*, London 2011.
60. Bhaskar A, Patel K, White A. Trigeminal Field Stimulation Using Percutaneous Electrical Nerve Stimulation (PENS) Therapy – preliminary case series of 8 patients. *Proceeding of 10th INS World Congress*, London 2011.
61. Khan EI, O'Keeffe D, Eller M, McFarland J, Walsh R, Gallagher H, Goroszenik T. Combined Spinal Cord and Subcutaneous Electrical Nerve Stimulation for the Treatment of Lower Extremity and Testicular Pain. *Proceedings of 6th INS World Congress*, 2003.
62. Racz GB, Ruiz-Lopez R. Radiofrequency procedures. *Pain Pract.* 2006 Mar;6(1):46-50.
63. Hooten WM, Martin DP, Huntoon MA. Radiofrequency neurotomy for low back pain: evidence-based procedural guidelines. *Pain Med.* 2005 Mar-Apr;6(2):129-38.
64. Soloman M, Mekhail MN, Mekhail N. Radiofrequency treatment in chronic pain. *Expert Rev Neurother.* 2010 Mar;10(3):469-74.
65. van Boxem K, van Eerd M, Brinkhuizen T, Patijn J, van Kleef M, van Zundert J. Radiofrequency and pulsed radiofrequency treatment of chronic pain syndromes: the available evidence. *Pain Pract.* 2008 Sep-Oct;8(5):385-93. Epub 2008 Aug 19. Review. Erratum in: *Pain Pract.* 2010 Mar-Apr;10(2):164. Brinkhuize, Tjinta [corrected to Brinkhuizen, Tjinta].
66. Markman JD, Philip A. Interventional approaches to pain management. *Anesthesiol Clin.* 2007 Dec;25(4):883-98, viii.
67. Golovac S. Radiofrequency neurolysis. *Neuroimaging Clin N Am.* 2010 May;20(2):203-14.
68. Lord SM, Bogduk N. Radiofrequency procedures in chronic pain. *Best Pract Res Clin Anaesthesiol.* 2002 Dec;16(4):597-617.
69. Bogduk N. Pulsed radiofrequency. *Pain Med.* 2006 Sep-Oct;7(5):396-407.
70. Protasoni M, Reguzzoni M, Sangiorgi S, Reverberi C, Borsani E, Rodella LF, Dario A, Tomei G, Dell'Orbo C. Pulsed radiofrequency effects on the lumbar ganglion of the rat dorsal root: a morphological light and transmission electron microscopy study at acute stage. *Eur Spine J.* 2009 Apr;18(4):473-8. Epub 2009 Jan 27
71. Tun K, Cemil B, Gurcay AG, Kaptanoglu E, Sargon MF, Tekdemir I, Comert A, Kanpolat Y. Ultrastructural Evaluation of Pulsed Radiofrequency and Conventional Radiofrequency lesions in rat sciatic nerve. *Surgical Neurol.* 2009 Nov; 72(5):496-500; discussion 501. Epub 2009 Mar 28

72. Hagiwara S, Iwasaka H, Takeshima N, Noguchi T. Mechanisms of analgesic action of pulsed radiofrequency on adjuvant-induced pain in the rat: roles of descending adrenergic and serotonergic systems. *Eur J Pain*. 2009 Mar; 13(3):249-52. Epub 2008 Jun 6
73. Takahashi N, Tasto JP, Ritter M, Ochiai N, Moriya H, Amiel D. Pain relief through an antinociceptive effect after radiofrequency application. *Am J Sports Med*. 2007 May;35(5):805-10. Epub 2007 Jan 31
74. Kanpolat Y, Savas A, Bekar A, Berk C. Percutaneous controlled radiofrequency trigeminal rhizotomy for the treatment of idiopathic trigeminal neuralgia: 25-year experience with 1,600 patients. *Neurosurgery*. 2001 Mar;48(3):524-32; discussion 532-4.
75. Emril DR, Ho KY. Treatment of trigeminal neuralgia: role of radiofrequency ablation. *J Pain Res*. 2010 Dec 12;3:249-54.
76. Erdine S, Ozyalcin NS, Cimen A, Celik M, Talu GK, Disci R. Comparison of pulsed radiofrequency with conventional radiofrequency in the treatment of idiopathic trigeminal neuralgia. *Eur J Pain*. 2007 Apr;11(3):309-13. Epub 2006 Jun 9.
77. Teixeira MJ, Siqueira SR, Almeida GM. Percutaneous radiofrequency rhizotomy and neurovascular decompression of the trigeminal nerve for the treatment of facial pain. *Arg Neuropsiquiatr*. 2006 Dec;64(4):983-9
78. van Kleef M, van Genderen WE, Narouze S, Nurmikko TJ, van Zundert J, Geurts JW, Mekhail N; World Institute of Medicine. 1. Trigeminal neuralgia. *Pain Pract*. 2009 Jul-Aug;9(4):252-9.
79. Madhusudan Reddy KR, Arivazhagan A, Chandramouli BA, Umamaheswara Rao GS. Multiple cranial nerve palsies following radiofrequency ablation for trigeminal neuralgia. *Br J Neurosurg*. 2008 Dec;22(6):781-3.
80. Savas A, Sayin M. Subarachnoid bleeding into the superior cerebellopontine cistern after radiofrequency trigeminal rhizotomy: case report. *Acta Neurochir(Wien)*. 2010 Mar;152(3):561-2. Epub 2009 Aug 11.
81. Ugur HC, Savas A, Elhan A, Kanpolat Y. Unanticipated complication of percutaneous radiofrequency trigeminal rhizotomy: rhinorrhea: report of three cases and a cadaver study. *Neurosurgery*. 2004 Jun;54(6):1522-4; discussion 1524-6.
82. Shah RV, Racz GB. Long-term relief of posttraumatic headache by sphenopalatine ganglion pulsed radiofrequency lesioning: a case report. *Arch Phys Med Rehabil*. 2004 Jun;85(6):1013-6.
83. Grégoire A, Clair C, Delabrousse E, Aubry R, Boulahdour Z, Kastler B. [CT-guided neurolysis of the sphenopalatine ganglion for management of refractory trigeminal neuralgia]. *J Radiol*. 2002 Sep;83(9 Pt 1):1082-4. French.
84. Meng FG, Wu CY, Liu YG, Liu L. Virtual reality imaging technique in percutaneous radiofrequency rhizotomy for intractable trigeminal neuralgia. *J Clin Neurosci*. 2009 Mar;16(3):449-51.
85. Rahman M, Richter EO, Osawa S, Rhoton AL Jr. Anatomic study of the infraorbital foramen for radiofrequency neurotomy of the infraorbital nerve. *Neurosurgery*. 2009 May;64(5 Suppl 2):423-7; discussion 427-8.
86. Vanelderen P, Rouwette T, De Vooght P, Puylaert M, Heylen R, Vissers K, VanZundert J. Pulsed radiofrequency for the treatment of occipital neuralgia: a prospective study with 6 months of follow-up. *Reg Anesth Pain Med*. 2010 Mar-Apr;35(2):148-51.
87. Navani A, Mahajan G, Kreis P, Fishman SM. A case of pulsed radiofrequency lesioning for occipital neuralgia. *Pain Med*. 2006 Sep-Oct;7(5): 453-6
88. Philip CN, Candido KD, Joseph NJ, Crystal GJ. Successful treatment of meralgia paresthetica with pulsed radiofrequency of the lateral femoral cutaneous nerve. *Pain Physician*. 2009 Sep-Oct;12(5):881-5.
89. Cohen SP, Foster A. Pulsed radiofrequency as a treatment for groin pain and orchialgia. *Urology*. 2003 Mar;61(3):645.
90. Rhame EE, Levey KA, Gharibo CG. Successful treatment of refractory pudendal neuralgia with pulsed radiofrequency. *Pain Physician*. 2009 May-Jun;12(3):633-8.
91. Mukubo Y, Sato K, Kawamata M, Masui. [Effective radiofrequency lesioning for obturator nerve neuropathy]. 2003 Sep;52(9):990-2. Japanese.
92. van Kleef M, Barendse GA, Dingemans WA, Wingen C, Lousberg R, de Lange S, Sluijter ME. Effects of producing a radiofrequency lesion adjacent to the dorsal root ganglion in patients with thoracic segmental pain. *Clin J Pain*. 1995 Dec;11(4):325-32.
93. Cohen SP, Sireci A, Wu CL, Larkin TM, Williams KA, Hurley RW. Pulsed radiofrequency of the dorsal root ganglia is superior to pharmacotherapy or pulsed radiofrequency of the intercostal nerves in the treatment of chronic postsurgical thoracic pain. *Pain Physician*. 2006 Jul;9(3):227-35.



94. Huang B, Zhou XY, Lu YP, Zhu ZF, Hou J, Sun JL, Xie GS, Zhu SM, Zheng HG. Zhonghua Yi Xue Za Zhi. [Selective percutaneous dorsal root ganglion radiofrequency thermocoagulation guided by CT scanning in treatment of post-herpetic neuralgia]. 2008 Apr 1;88(13):885-8. Chinese.
95. Uchida K. Radiofrequency treatment of the thoracic paravertebral nerve combined with glucocorticoid for refractory neuropathic pain following breast cancer surgery. *Pain Physician*. 2009 Jul-Aug; 12(4):E277-83.
96. Restrepo-Garces CE, Marinov A, McHardy P, Faclier G, Avila A. Pulsed radiofrequency under ultrasound guidance for persistent stump-neuroma pain. *Pain Pract*. 2011 Jan-Feb;11(1):98-102. doi: 10.1111/j.1533-2500.2010.00398.x.
97. Wilkes D, Ganceres N, Solanki D, Hayes M. Pulsed radiofrequency treatment of lower extremity phantom limb pain. *Clin J Pain*. 2008 Oct;24(8):736-9.
98. West M, Wu H. Pulsed radiofrequency ablation for residual and phantom limb pain: a case series. *Pain Pract*. 2010 Sep-Oct;10(5):485-91.
99. Tamimi MA, McCeney MH, Krutsch J. A case series of pulsed radiofrequency treatment of myofascial trigger points and scar neuromas. *Pain Med*. 2009 Sep;10(6):1140-3. Epub 2009 Jul 6.
100. Shabat S, Pevsner Y, Folman Y, Gepstein R. Pulsed radiofrequency in the treatment of patients with chronic neuropathic spinal pain. *Minim Invasive Neurosurg*. 2006 Jun;49(3):147-9.
101. Cozzarelli J, Sollitto RJ, Thapar J, Caponigro J. A 12-year long-term retrospective analysis of the use of radiofrequency nerve ablation for the treatment of neurogenic heel pain. *Foot Ankle Spec*. 2010 Dec; 3(6):338-46. Epub 2010 Sep 3
102. Liden B, Simmons M, Landsman AS. A retrospective analysis of 22 patients treated with percutaneous radiofrequency nerve ablation for prolonged moderate to severe heel pain associated with plantar fasciitis *J Foot Ankle Surg*. 2009 Nov-Dec;48(6):642-7. Epub 2009 Jul 5.
103. Straube S, Derry S, Moore RA, McQuay HJ. Cervico-thoracic or lumbar sympathectomy for neuropathic pain and complex regional pain syndrome. *Cochrane Database Syst Rev*. 2010 Jul 7;(7):CD002918.
104. Forouzanfar T, van Kleef M, Weber WE. Radiofrequency lesions of the stellate ganglion in chronic pain syndromes: retrospective analysis of clinical efficacy in 86 patients. *Clin J Pain*. 2000 Jun;16(2):164-8.
105. Akkoc Y, Uyar M, Oncu J, Ozcan Z, Durmaz B. Complex regional pain syndrome in a patient with spinal cord Injury: management with pulsed radiofrequency lumbar sympathectomy. *Spinal Cord*. 2008 Jan;46(1):82-4. Epub 2007 Jul 31.
106. Mailis A, Furlan A. Sympathectomy for neuropathic pain. *Cochrane Database Syst Rev*. 2003;(2): CD002918. Review. Update in: *Cochrane Database Syst Rev*. 2010;(7):CD002918.